

EGU2020-5799

<https://doi.org/10.5194/egusphere-egu2020-5799>

EGU General Assembly 2020

© Author(s) 2020. This work is distributed under the Creative Commons Attribution 4.0 License.



The impact of the Messinian Salinity Crisis on marine biota

Konstantina Agiadi¹, Niklas Hohmann², Giorgio Carnevale³, Elsa Gliozzi⁴, Constanza Faranda⁴, Francesca Lozar³, Mathias Harzhauser⁵, George Iliopoulos⁶, Antonio Caruso⁷, George Kontakiotis¹, Marco Taviani⁸, Alan Maria Mancini³, Enrico Borghi⁹, Ildefonso Bajo Campos¹⁰, Pierre Moissette¹, Danae Thivaoui¹, Stergios Zarkogiannis¹, Eva Besiou¹, Daniel Garcia-Castellanos¹¹, and Angelo Camerlenghi¹²

¹Department of Historical Geology and Paleontology, National and Kapodistrian University of Athens, Athens, Greece
(kagiadi@geol.uoa.gr)

²Friedrich-Alexander-Universität GeoZentrum Nordbayern, Erlangen, Germany

³Department of Earth Sciences, University of Torino, Torino, Italy

⁴Dipartimento di Scienze-Sezione Scienze Geologiche, Università Roma 3, Rome, Italy

⁵Natural History Museum, Vienna, Austria

⁶Faculty of Geology, University of Patras, Rio, Greece

⁷Dipartimento di Scienze della Terra e del Mare, Università degli studi di Palermo, Palermo, Italy

⁸University of Bologna, Institute of Marine Science - National Research Council ISMAR-CNR, Bologna, Italy

⁹Società Reggiana di Scienza Naturali

¹⁰Sección de Paleontología, Museo de Alcalá de Guadaíra, Seville, Spain

¹¹ICTJA-CSIC, Barcelona, Spain

¹²OGS Istituto Nazionale di Oceanografia e di Geofisica Sperimentale, Trieste, Italy

The Messinian Salinity Crisis (MSC) was the greatest paleoenvironmental perturbation the Mediterranean has ever seen. The literature is abundant in hypotheses on the repercussions of the MSC on organisms. However, all these are based on incomplete and still uncertain scenarios about the MSC evolution, as well as on the assumption that such a paleoenvironmental perturbation must have completely reset marine biota. Having prevailed for many decades now, this assumption has leaked from paleontology and geosciences to biological sciences, with numerous studies taking this scenario for granted instead of using it as a starting hypothesis to be tested. Here, we review and revise the marine fossil record across the Mediterranean from the Tortonian until the Zanclean to follow the current rules of nomenclature, correct misidentifications, and control for stratigraphic misplacements. We examine the composition of marine faunas, both taxonomically and considering the function of each group in the marine ecosystem and the transfer of energy through the marine food web. Specifically, we investigate the following functional groups: 1) primary producers, 2) secondary producers, 3) primary consumers, 4) secondary consumers, and 5) top predators. Our study includes sea grasses, phytoplankton, corals, benthic and planktonic foraminifera, bivalves, gastropods, brachiopods, echinoids, bryozoans, fishes, ostracods, and marine mammals. We calculate biodiversity indexes to provide independent evidence quantifying to what degree the marine fauna underwent:

- A drop of overall regional biodiversity of the Mediterranean due to environmental stress during the Messinian.
- A taxonomic and functional change between the Tortonian, Messinian, and the Zanclean, that is before and after the MSC, as well as during the precursor events to that actual crisis taking place after the Tortonian/Messinian boundary.
- The onset of the present-day west-to-east decreasing gradient in species richness, which has been related to the sea temperature and productivity gradients and the distance from the Gibraltar connection to the Atlantic.